

An Alternative Parametrization of the Pion Form Factor and the Mass and Width of $\rho(770)^*$

Peter Lichard^{1,2} and Martin Voják¹

¹*Institute of Physics, Silesian University in Opava, Bezručovo nám. 13, 746 01 Opava, Czech Republic*

²*Institute of Experimental and Applied Physics, Czech Technical University, Horská 3/a, 120 00 Prague, Czech Republic*

(Dated: February 2, 2008)

In order to reveal possible mass shifts of the vector mesons in a dense strongly interacting medium presumably created in high energy heavy ion collisions it is necessary to know their free masses reliably. The $\rho(770)$ mass quoted in the last two editions of the Review of Particle Physics is significantly larger than the values quoted in previous editions. The new value is mostly influenced by the results of recent experiments CMD-2 and SND at the VEPP-2M e^+e^- collider at Novosibirsk. We show that the values of the mass and width of the $\rho(770)$ meson measured in the $e^+e^- \rightarrow \pi^+\pi^-$ annihilation depend crucially on the parametrization used for the pion form factor. We propose a parametrization of the $\rho(770)$ contribution to the pion form factor based on the running mass calculated from a single-subtracted dispersion relation and compare it with the parametrization based on the formula of Gounaris and Sakurai used recently by the CMD-2 collaboration. We show that our parametrization gives equally good or better fits when applied to the data of the CMD-2, SND, and KLOE collaborations, but yields much smaller values of the $\rho(770)$ mass, consistent with the photoproduction and hadronic reactions results. Our fit to the KLOE data becomes exceptionally excellent (confidence level of 99.88%) if an energy shift of about 2 MeV in the $\rho\omega$ region is allowed.

I. INTRODUCTION

The vector meson $\rho(770)$ plays an important role in many phenomena of particle and nuclear physics. Possible modification of its mass and width in dense strongly interacting systems presumably created in high energy heavy ion collisions is now widely discussed [1–11]. To study this effect quantitatively, reliable information about the “vacuum” values of these parameters is desirable.

Inspecting recent editions of the Review of Particle Physics [12–19] we find: (i) whereas the values of $\rho(770)$ mass quoted until 2002 are compatible within experimen-

TABLE II: Mass of ρ measured in different processes, selected and averaged by PDG in 2006 [19]

	m_{ρ^0} (MeV)	$m_{\rho^\pm}$ (MeV)
$e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0$	775.5 ± 0.4	775.4 ± 0.4^a
Photoproduction	768.5 ± 1.1	/
Hadronic reactions	769.0 ± 0.9	766.5 ± 1.1

^aIncludes also τ decay results

tal errors, an abrupt rise appears in 2004 (see Table I); (ii) the ρ^0 mass average calculated from the e^+e^- annihilation experiments is much higher than the averages calculated from hadronic reactions and photoproduction. The figures taken from the 2006 Particle Data Group (PDG) tables are shown in Table II.

The discrepancy between the e^+e^- annihilation measurements of the ρ^0 mass and the results of photoproduction and hadronic experiments is very troublesome. We know of no single theoretical prediction or explanation of such a phenomenon.

In the literature one can find many articles that dealt with the determination of the $\rho(770)$ parameters from the experimental data on the pion form factor [20–30]. Their results on the ρ^0 mass are smaller, in some cases substantially, than the value preferred now by the PDG. It is not clear what has led the PDG to not using them “for averages, fits, limits, etc.” [19]. The PDG average builds on five experiments, three of which explored the e^+e^- annihilation into a pair of charged pions. Two of those three experiments utilized the Gounaris-Sakurai (GS) parametrization [31] when fitting the data.

As we discuss in detail in the next section, the valid-

TABLE I: Mass and width of $\rho(770)$ according to PDG

year	m_ρ (MeV)	Γ_ρ (MeV)
1992 [12]	768.1 ± 0.5	151.5 ± 1.2
1994 [13]	769.9 ± 0.8	151.2 ± 1.2
1996 [14]	768.5 ± 0.6	150.7 ± 1.2
1998 [15]	770.0 ± 0.8	150.7 ± 1.1
2000 [16]	769.3 ± 0.8	150.2 ± 0.8
2002 [17]	771.1 ± 0.9	149.2 ± 0.7
2004 [18]	775.8 ± 0.5	150.3 ± 1.6
2006 [19]	775.5 ± 0.4	149.4 ± 1.0

*This paper is dedicated to the late Julia Thompson, who drew the attention of one of us (P. L.) to the experimental program of the Budker Institute of Nuclear Physics at Novosibirsk.

ity of the GS formula is limited because it ignores two important effects. Namely, the opening of additional decay channels at higher energies and the structure effects in the hadronic vertices manifesting themselves as the momentum dependent strong form factors. These effects directly influence the imaginary part in the denominator of the GS formula. This itself would not have serious implications for the analysis of the form factor data, which are usually restricted to a narrow interval around the nominal ρ mass. But because of the close connection between the denominator's real and imaginary part implied by the analyticity, the real part is also influenced, even at the energies around the nominal ρ mass, which are important for data fitting.

Already in the pioneer era of the ρ -meson physics, when the ρ mass and width were determined mainly by analyzing the dipion mass distribution in the reaction $\pi p \rightarrow \pi\pi N$, Pišút and Roos [32] showed that the discrepancies among the results of different experimental groups are caused by different parametrizations these groups used. They also discussed the discrepancy between the hadronic measurements and the results of the first e^+e^- annihilation into two pions experiment [33]. From that time the parametrization dependence, mainly in connection with the pion form factor parametrization, has been discussed in many papers, see, e.g., Refs. [24, 26, 28, 30, 34, 35].

The aim of this paper is to show that the uncertainties connected with the choice of the form factor parametrization induce much larger errors of the ρ resonance parameters than the statistical and systematic errors of the data. For that purpose we fit three different data sets with two different parametrizations and compare the resulting masses and width of the ρ^0 meson. One of the parametrizations is that used recently by the CMD-2 collaboration [36, 37]. Its principal element is the GS formula. The other parametrization [38] is based on a single-subtraction dispersion relation fed by the imaginary part taking into account several decay channels which open when the ρ -resonance goes “off-mass-shell” towards higher masses. All decay widths are calculated taking into account the strong form factors.

The paper is organized as follows. In Sec. II we describe the two versions of the ρ^0 resonance term in the pion form factor. The parametrization of the subleading contributions coming from the $\rho(1450)$, $\omega(782)$, and $\phi(1020)$ is presented in Sec. III. The results of fitting the CMD-2 [37], SND [39], and KLOE [40] data sets by the two parametrizations are shown and discussed in Sec. IV. Section V contains summary and concluding remarks.

II. TWO PARAMETRIZATIONS OF THE $\rho(770)$ CONTRIBUTION TO THE PION FORM FACTOR

We write the contribution of the intermediate state with the ρ^0 meson to the pion electromagnetic form factor

in the following form

$$F_\rho(s) = \frac{M_\rho^2(0)}{M_\rho^2(s) - s - im_\rho\Gamma_\rho(s)}, \quad (1)$$

where the running mass squared $M_\rho^2(s)$ is a real function satisfying the conditions

$$M_\rho^2(m_\rho^2) = m_\rho^2, \quad (2)$$

and

$$\left. \frac{dM_\rho^2(s)}{ds} \right|_{s=m_\rho^2} = 0, \quad (3)$$

m_ρ is the nominal mass of the ρ meson, and the real function $\Gamma_\rho(s)$ is the total decay width of a ρ^0 meson with mass $\sqrt{s}$. It follows from microcausality that the pion form factor $F(s)$ is an analytic function in the whole complex s -plane with a cut running from $s = 4m_\pi^2$ to infinity, satisfying the condition $F(s^*) = F^*(s)$, and growing no faster than a finite power of s for $|s| \rightarrow \infty$ (in QCD, $|F(s)|$ even decreases [41]). These properties of $F(s)$ imply the analytic properties of the denominator in Eq. (1).

A. Gounaris-Sakurai formula

Gounaris and Sakurai [31] assumed a low-energy parametrization of the phase shift in the P_{11} wave of the $\pi\pi$ elastic scattering inspired by work of Chew and Mandelstam [42] and used the relation between the pion form factor and the $\pi\pi$ elastic scattering amplitude in the P_{11} channel [43]. After defining the resonance parameters in terms of the P_{11} phase shift they obtained a parametrization of the pion form factor. Below, we write it in a little different form, adopting partly the notation of Ref. [36] and referring to our Eq. (1). Firstly, on the upper branch of the cut we have

$$\Gamma_\rho(s) = \Gamma_\rho \frac{m_\rho}{\sqrt{s}} \left(\frac{s - 4m_\pi^2}{m_\rho^2 - 4m_\pi^2} \right)^{3/2}, \quad (4)$$

where $\Gamma_\rho = \Gamma_\rho(m_\rho^2)$ is the nominal width of the $\rho(770)$ meson, whereas $\Gamma_\rho(s) \equiv 0$ for $s \leq 4m_\pi^2$. Secondly,

$$M_\rho^2(s) = m_\rho^2 + f(s), \quad (5)$$

where

$$f(s) = \frac{2\Gamma_\rho m_\rho^2}{(m_\rho^2 - 4m_\pi^2)^{3/2}} \left\{ (s - 4m_\pi^2)[h(s) - h(m_\rho^2)] + (m_\rho^2 - 4m_\pi^2)h'(m_\rho^2)(m_\rho^2 - s) \right\}, \quad (6)$$

$$h(s) = \begin{cases} \frac{1}{\pi} \sqrt{\frac{s-4m_\pi^2}{s}} \ln \frac{\sqrt{s} + \sqrt{s-4m_\pi^2}}{2m_\pi}, & s \geq 4m_\pi^2, \\ \sqrt{\frac{4m_\pi^2-s}{4s}} \left[1 - \frac{2}{\pi} \arctan \sqrt{\frac{4m_\pi^2-s}{s}} \right], & 0 < s < 4m_\pi^2, \\ \frac{1}{\pi}, & s = 0, \\ \frac{1}{\pi} \sqrt{\frac{s-4m_\pi^2}{s}} \ln \frac{\sqrt{-s} + \sqrt{4m_\pi^2-s}}{2m_\pi}, & s < 0. \end{cases}$$

To get more insight into the physics behind the GS parametrization let us utilize that $M_\rho^2(s)$ and $\Gamma_\rho(s)$, which are related to the ρ -meson contribution to the pion form factor by Eq. (1), should be the boundary values of the real and imaginary part, respectively, of a function analytic in the cut s -plane. It is easy to check that the quantities (4) and (5) are related through the double-subtracted dispersion relation

$$M_\rho^2(s) = A_0 + A_1 s - \frac{s^2}{\pi} \mathcal{P} \int_{4m_\pi^2}^{\infty} \frac{m_\rho \Gamma_\rho(s')}{s'^2(s' - s)} ds'. \quad (7)$$

The appearance of the two subtraction constants A_0 and A_1 guarantees that the conditions (2) and (3) can be satisfied. Of course, the relation (7) is not specific for the GS parametrization, but is more general. It enables us to find the running mass squared $M_\rho^2(s)$ corresponding to any reasonable (i.e., not growing too fast) choice of the total width $\Gamma_\rho(s)$. It is important to realize that the values of $M_\rho^2(s)$ at any s , in particular those for low $s < 1 \text{ GeV}^2$, which are used in the form factor fitting, are given by the values of $\Gamma_\rho(s)$ at all s . To get reliable $M_\rho^2(s)$ it is therefore essential to have $\Gamma_\rho(s)$ that is physically sound not only in the s -region vital for the fitting, but also beyond it.

The GS parametrization of the $\rho(770)$ contribution to the pion electromagnetic form factor ignores possible structure effects in the $\rho^0 \pi^+ \pi^-$ vertex (the strong form factor). They were taken into account by Vaughn and Wali (VW) [44] in the form of a phenomenological cutoff function. When that function is set to unity, the VW parametrization reduces to the GS one. The VW approach has not become as popular as that of GS, although it is closer to reality, because the outcome of dispersion relation (7) cannot be expressed in an analytic form, similar to (6).

A different attempt to bring the GS-type formulas closer to reality was done by the authors of Ref. [30]. They included also the contributions from the $K^+ K^-$ and $K^0 \bar{K}^0$ states to the imaginary part of the denominator of the pion form factor (1) and obtained its real part in an analytic form using the dispersion relation. No strong form factor was included. Also this improvement of the GS formula passed unnoticed by the experimentalists fitting their data.

We should also note that the original derivation of the GS formula brings a one-to-one correspondence between the P_{11} scattering length and effective radius on one side, and the $\rho(770)$ resonance position and width on the other. Using the GS formula for the contribution of higher resonances [$\rho(1450)$, $\rho(1700)$] to the pion form factor, as in [36, 37, 45, 46], thus contradicts the spirit of the GS derivation as it implies incorrect values of the P_{11} scattering length and effective radius. In the language of dispersion relation (7) it corresponds to the assumption that the two-pion mode is the only decay mode at any $\sqrt{s}$. This is unrealistic because already for the $\rho(1450)$ with nominal mass there are several other modes allowed by the energy-momentum, angular momentum, and internal quantum numbers conservation.

B. Our running mass approach

We build our alternative parametrization on the recent work by one of us [38], to which the reader is referred for technical details. The numerical values of the constants used in [38] are updated using the 2002 edition of the Review of Particle Physics [17].

There are four items in which our parametrization differs from the GS one:

(i) We consider several contributions to the total decay width of the ρ meson, which become operational when its mass $\sqrt{s}$ exceeds respective thresholds

$$\begin{aligned} \Gamma_\rho(s) = & \Gamma_{\rho \rightarrow \pi\pi}(s) + \Gamma_{\rho \rightarrow \omega\pi}(s) \\ & + \Gamma_{\rho \rightarrow K\bar{K}}(s) + \Gamma_{\rho \rightarrow \eta\pi\pi}(s). \end{aligned} \quad (8)$$

(ii) The partial decay width of the two-pion decay, which provides the essential contribution to (8) is calculated from the standard interaction Lagrangian containing the first derivative of the pion field. The result is

$$\Gamma_{\rho \rightarrow \pi\pi}(s) = \Gamma_\rho \frac{m_\rho^2}{s} \left(\frac{s - 4m_\pi^2}{m_\rho^2 - 4m_\pi^2} \right)^{3/2}. \quad (9)$$

(iii) The strong form factors are introduced into all purely hadronic vertices. They were taken from the flux-tube-breaking model of Kokoski and Isgur [47]. This leads to the modification of the partial decay widths composing (8) and calculated originally as if the participating mesons were elementary quanta. For example, the two-pion width (9) is modified by the following multiplication factor

$$\exp \left\{ \frac{m_\rho^2 - s}{24\beta^2} \right\},$$

where the value $\beta = 0.4$ was taken from the Kokoski and Isgur [47] analysis.

(iv) As now the the total width (8) does not rise so fast as in the GS approach, we may write a once-subtracted




FIG. 1: Comparison of the running mass squared of ρ -meson in the Gounaris-Sakurai and our approach

dispersion relation

$$M_\rho^2(s) = B_0 - \frac{s}{\pi} \mathcal{P} \int_{4m_\pi^2}^{\infty} \frac{m_\rho \Gamma_\rho(s')}{s'(s' - s)} ds'. \quad (10)$$

The subtraction constant B_0 is chosen to ensure that the condition $M_\rho^2(m_\rho^2) = m_\rho^2$ (2) is satisfied. Now it is not possible to satisfy the condition on derivative (3) automatically. We consider this an advantage, because Eq. (3) can thus serve as a test that the energy dependent total decay width (8) was chosen correctly. For details, see Ref. [38]. Let us note that our choice of the subtraction point $s = 0$ differs from $s = m_\rho^2$, used in Refs. [38, 48]. The small disadvantage lying in the necessity to evaluate the subtraction constant B_0 is generously compensated by much faster numerical evaluation of the dispersion integral. In the following, we are going to vary m_ρ and Γ_ρ in a search for the best fit to the experimental data. It means that we will have to calculate the running mass (10) in the experimental points again and again. The computation speed is therefore a decisive factor.

The comparison of the basic quantities of the GS and our approach is shown in Fig. 1 (running mass) and Fig. 2 (total width). The values $m_\rho = 771.1$ MeV/ c^2 and $\Gamma_\rho = 149.2$ MeV [17] were chosen for this purpose.

III. OTHER CONTRIBUTIONS TO THE PION FORM FACTOR

The parametrization of the pion electromagnetic form factor should contain, in addition to the dominant contri-




FIG. 2: Comparison of the energy dependent width of ρ -meson in the Gounaris-Sakurai and our approach

bution from the $\rho(770)^0$, also the contribution from other resonances. In the energy range of the $e^+e^- \rightarrow \pi^+\pi^-$ experiments we are interested in, they include $\rho' \equiv \rho(1450)$, $\omega(782)$, and perhaps also $\phi(1020)$. We write the pion form factor in the form

$$F(s) = \left\{ F_\rho(s) \left[1 + \delta \frac{s}{m_\omega^2} F_\omega(s) \right] + \beta F_{\rho'}(s) + \gamma F_\phi(s) \right\} \times (1 + \beta + \gamma)^{-1}, \quad (11)$$

where not only $F(s)$ but also the individual contributions to it are normalized to unity at $s = 0$. Parameters β , γ , and δ are complex and should be determined, together with m_ρ and Γ_ρ by minimizing χ^2 constructed from the data and parametrizations in the standard way. Equation (11) is inspired by Eq. (8) from [36] and reduces to it for $\gamma = 0$. We will use it in conjunction with both GS and our running mass parametrizations. In one case (CMD-2 data) allowing non-vanishing γ did not bring a considerable drop in χ^2 . But for the other two data sets (SND, KLOE) the ϕ contribution improved the fits.

The CMD-2 collaboration [36, 37] used the GS parametrization also for the $\rho(1450)$ contribution and we will do the same when applying their formalism. But when using our parametrization, we utilize the fixed-mass formula

$$F_{\rho'}(s) = \frac{m_{\rho'}^2}{m_{\rho'}^2 - s - im_{\rho'}\Gamma_{\rho'}(s)} \quad (12)$$

with several options for the total width of ρ' :

1. $\Gamma_{\rho'}(s) \equiv \Gamma_{\rho(1450)}$;

2. Eq. (4) adapted for $\rho(1450)$;
3. $\Gamma_{\rho'}(s) \equiv 0$;
4. Eq. (9) adapted for $\rho(1450)$.

The options 2 and 4 are equivalent to the assumption that the only decay mode of $\rho(1450)$ is the dipion one. The option 3 mimics the situation when $\rho(1450)$ disintegrates only through the modes with higher thresholds, the decay widths of which become negligible below $s = 1 \text{ GeV}^2$ mainly due to the shrinking phase-space. This is in conformity with what was known already a long time ago, namely that the ρ' resonance is clearly visible in the e^+e^- annihilation into four pions [49] whereas its appearance in the two-pion channel is less pronounced, see, e.g., Fig. 2 in [46].

The results of the fits (χ^2 , $m_{\rho(770)}$, $\Gamma_{\rho(770)}$) showed only marginal differences among various options, with option 3 giving slightly better χ^2 than the other options. The results of our approach presented below were obtained using just that option.

As to the contribution of the $\omega(782)$ to the pion electromagnetic form factor, we use

$$F_\omega(s) = \frac{m_\omega^2}{m_\omega^2 - s - im_\omega \Gamma_\omega(s)}. \quad (13)$$

The following options are considered:

1. $\Gamma_\omega(s) \equiv \Gamma_{\omega(782)}$ above the $s = 9m_\pi^2$ threshold;
2. $\Gamma_\omega(s) = \Gamma_{\omega \rightarrow 3\pi}(s)$;
3. same as 2, but including strong form factors;
4. $\Gamma_\omega(s) = \Gamma_{\omega \rightarrow 3\pi}(s) + \Gamma_{\omega \rightarrow \pi\gamma}(s) + \Gamma_{\omega \rightarrow \pi^+\pi^-}(s)$;
5. same as 4, but including strong form factors.

Following [36], the option 1 was used in conjunction with the GS parametrization of the $\rho(770)$ and $\rho(1450)$ contributions. The other options were explored as a part of our approach, but did not exhibit notable differences in the quality of the fits.

In the case of the SND data, the contribution of the ρ - ϕ interference is present in a statistically significant way, as judged from the χ^2 decrease, with both parametrizations (GS, ours) which we explore. For data of the KLOE collaboration, the situation is more complicated. The details are reported in Sec. IV C. Our parametrization of the $\phi(1020)$ contribution to the pion form factor is

$$F_\phi(s) = \frac{m_\phi^2}{m_\phi^2 - s - im_\phi \Gamma_\phi(s)}. \quad (14)$$

We consider the following options:

1. $\Gamma_\phi(s) \equiv \Gamma_{\phi(1020)}$ above the $s = 4m_K^2$ threshold;
2. $\Gamma_\phi(s) = \Gamma_{\phi \rightarrow K\bar{K}}(s)$;
3. same as 2, but including strong form factor;

$$4. \Gamma_\phi(s) = \Gamma_{\phi \rightarrow K\bar{K}}(s) + \Gamma_{\phi \rightarrow 3\pi}(s) + \Gamma_{\phi \rightarrow \eta\gamma}(s);$$

5. same as 4, but including strong form factors.

Given the narrow width of the ϕ resonance, the high thresholds of its important decay modes, and the distance of the experimental energies from the ϕ mass, the refinements 2–5 did not change the results noticeably in comparison with the simplest option 1.

IV. RESULTS

A. Using the CMD-2 experiment data

In 2002, the CMD-2 Collaboration published [36] their results of the precise measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section for 43 c.m. energies and gave also the values of the pion electromagnetic form factor determined from them. After an error in the computer code was found, they published the updated values of the pion form factor [37], which we are going to explore here. The original paper [36] remains important as it contains details about the GS based parametrization, which the collaboration used in fitting the form factor data.

We first tested our computer code using the uncorrected CMD-2 data because in Ref. [36] a more complete information is provided than in the later paper [37]. Our results agreed with the values in Table 3 [36]. Then we used the corrected pion form factor data [37] and got the results shown in Tab. III. The mass and width of the ρ meson we obtained differ only very slightly from the CMD-2 values, differences are much smaller than the errors quoted in [37]. Our errors shown in Tab. III and elsewhere are only the statistical ones in the MINUIT [50] sense.

Our running mass parametrization gives a little better fit to the CMD-2 data than the GS one. The results are shown in the last column of Tab. III. The $\rho(770)$ mass and width we got are much lower than the corresponding quantities obtained by the GS parametrization.

It is also interesting to compare the mean square radius of the charged pion, which is related to the derivative of the form factor at $s = 0$ [51], with the experimental value $\langle r_\pi^2 \rangle = (0.439 \pm 0.008) \text{ fm}^2$ [52].

The graphical comparison of our parametrization with the CMD-2 data is shown in Fig. 3.

B. Using the SND experiment data

In 2005, the results were reported [53] of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section measurement with the spherical neutral detector (SND) at the VEPP-2M collider in Novosibirsk. Later on, an error in software was found and the data corrected [39]. We use the data on pion form factor from their Table I. The experimental value of the charged pion mean square radius [52] is added to the fitted data

TABLE III: Mass and width of $\rho(770)$ and the mean-square radius (msr) of the pion from fitting the CMD-2 data. The experimental value of the msr is $0.439 \pm 0.008 \text{ fm}^2$ [52].

	GS	ours
χ^2/NDF	35.16/37	32.18/37
$m_\rho \text{ (MeV}/c^2)$	775.3 ± 1.1	767.08 ± 0.83
$\Gamma_\rho \text{ (MeV)}$	143.2 ± 2.5	136.1 ± 2.2
$\langle r_\pi^2 \rangle \text{ (fm}^2)$	0.416 ± 0.013	0.434 ± 0.012




FIG. 3: Comparison of our parametrization of the pion form factor with the CMD-2 data [37]

set. The comparison of the GS and our parametrization (both include also the contribution from the ϕ meson) is reported on in Table IV. We also reproduce there the results of the updated [39] fit by the SND collaboration. The formulas used in their fitting are described in [53] together with three choices of the ρ' and ρ'' parameters. Unfortunately, in the updated paper the provided information is scarce. It is not said which choice was used and what was the χ^2 of the fit. Nevertheless, there is a good compatibility in both m_ρ and Γ_ρ between the GS parametrization and that used by the SND collaboration.

The GS and our parametrization provide equally good fits, but the mass and width resulting from the latter are lower than and incompatible with those from the former.

C. Using the KLOE experiment data

The data on the cross section $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma)$ at the collision energy $W = 1.02 \text{ GeV}$ measured with the KLOE detector at the electron-positron collider DAΦNE

TABLE IV: Mass and width of $\rho(770)$ from fitting the SND data.

	GS	ours	SND fit [39]
χ^2/NDF	39.94/38	39.80/38	not given
$m_\rho \text{ (MeV}/c^2)$	774.60 ± 0.52	764.23 ± 0.43	$774.6 \pm 0.4 \pm 0.5$
$\Gamma_\rho \text{ (MeV)}$	147.3 ± 1.0	140.29 ± 0.85	$146.1 \pm 0.8 \pm 1.5$




FIG. 4: Comparison of our parametrization of the pion form factor with the SND data [39]

in Frascati were used in Ref. [40] to extract the square of the charged pion form factor $|F_\pi(s)|^2$ by the radiative return method [54]. Here, $s < W^2$ denotes the square of the invariant mass of the $\pi^+\pi^-$ system. No attempt was made by the authors of [40] to fit their form factor in order to get the ρ meson parameters. We have performed the fit using both GS and our parametrizations. The results are summarized in Table V. The χ^2 of both fits is poor. Nevertheless, the χ^2 of our parametrization is about half that of the GS parametrization. An interesting observation is that not only our parametrization, but also the GS one gives a $\rho(770)$ mass substantially smaller than both Novosibirsk experiments. As before, our parametrization yields smaller values of both the mass and width than GS. An attempt to improve the quality of the fits by including the ρ - ϕ interference somewhat lowered the χ^2 for both parametrizations, but not enough to get the confidence level into an acceptable range.

Figure 5 shows that the worst discrepancy of our parametrization with the data is located in the ρ - ω region. The abrupt fall, which is present in Figs. 3 and 4, is missing. A more detailed analysis of the latter figures

TABLE V: Mass and width of $\rho(770)$ from fitting the KLOE data.

	GS	ours
χ^2/NDF	367.8/54	192.6/54
m_ρ (MeV/ c^2)	769.22 ± 0.32	761.39 ± 0.25
Γ_ρ (MeV)	144.71 ± 0.56	140.07 ± 0.66




FIG. 5: Comparison of our parametrization of the pion form factor with the KLOE data [40].

showed that the steep descent is caused by the ρ - ω interference term, which is very sensitive to the exact position of the (very narrow) ω resonance. The absence of the interference term in Fig. 5 suggests that the location of the ω resonance in the KLOE data is different from the nominal one, which we assumed when doing the fit with both GS and our parametrization. It would mean that the energy scale in the KLOE data contains a systematic shift, at least in the ρ - ω region. The KLOE collaboration utilized the radiative return method [54]. This means that the invariant mass of the $\pi^+\pi^-\gamma$ system (denoted as W in [40]) is fixed by the energy of the colliding e^+ and e^- . But the invariant mass $\sqrt{s}$ of the $\pi^+\pi^-$ subsystem has to be calculated from the measured momenta of the outgoing particles and is therefore known less precisely.

To scrutinize the possibility of a shift in experimental values of $\sqrt{s}$, we repeated the fits allowing the ω mass also to vary. Both the GS and our parametrization indicate that the best fit is obtained if the ω mass is smaller than the nominal one by about 2 MeV, see Table VI. The drop of χ^2 in both parametrizations justifies including the ω mass among free parameters, compare Table VI with Table V. But the quality of the fit is very




FIG. 6: Comparison of our parametrization of the pion form factor with the KLOE data [40] after decreasing the $\omega(782)$ mass parameter by 1.93 MeV.

TABLE VI: Fitting the KLOE data, the $\omega(782)$ mass allowed to vary.

	GS	ours
χ^2/NDF	202.6/53	29.58/53
m_ω (MeV/ c^2)	780.52 ± 0.33	780.72 ± 0.15
Δm_ω (MeV/ c^2) ^a	2.13 ± 0.35	1.93 ± 0.19
m_ρ (MeV/ c^2)	769.45 ± 0.25	761.51 ± 0.14
m_ρ (MeV/ c^2) ^b	771.58 ± 0.60	763.44 ± 0.33
Γ_ρ (MeV)	146.39 ± 0.51	142.56 ± 0.41

^aDifference from $m_\omega = (782.65 \pm 0.12)$ MeV [19]

^bThe value resulting from the fit was increased by Δm_ω

different for the two parametrizations. While the value of $\chi^2 = 202.6$ for the GS parametrization is still very high (almost four times the number of degrees of freedom), our parametrization yields $\chi^2 = 29.6$, which means, considering $NDF = 53$, the confidence level of 0.9962. Such perfect fits are rarely encountered in the high energy physics. The improvement of the fit is clearly seen by naked eye if one compares the new parametrization (Fig. 6) with the old one (Fig. 5).

Assuming that our analysis really signifies that a systematic shift of $\sqrt{s}$ is present in the data, we used the difference Δm_ω between the nominal mass of the $\omega(782)$ and that resulting from our fit to get the “corrected” ρ -meson mass, see Table VI.

Unfortunately, our interpretation of the mass shift in the ρ - ω mass region as an experimental effect is difficult to be explained in terms of a genuine miscalibration of

TABLE VII: As Tab. VI, but including the ρ - ϕ interference.

	GS	ours
χ^2/NDF	58.35/51	25.67/51
m_ω (MeV/ c^2)	780.15 ± 0.17	780.71 ± 0.15
Δm_ω (MeV/ c^2) ^a	2.50 ± 0.21	1.94 ± 0.19
m_ρ (MeV/ c^2)	770.29 ± 0.16	761.54 ± 0.16
m_ρ (MeV/ c^2) ^b	772.79 ± 0.35	763.48 ± 0.33
Γ_ρ (MeV)	151.53 ± 0.55	142.78 ± 0.46

^aDifference from $m_\omega = (782.65 \pm 0.12)$ MeV [19]^bThe value resulting from the fit was increased by Δm_ω

TABLE VIII: Confidence level of the GS and our fits to various data sets

	GS	ours
CMD-2	0.5555	0.6943
SND	0.3840	0.3899
KLOE	$< 10^{-30}$	2×10^{-17}
KLOE ^a	2×10^{-19}	0.9962
KLOE ^b	0.2234	0.9988

^aWith the ω mass shifted^bWith the ω mass shifted and ϕ contribution included

the KLOE drift chamber [55].

In an effort to reveal the origin of the mysterious mass shift we considered the possibility that it was an artifact of a wrong parametrization. We combined our parametrization of the ρ^0 contribution (1) to the pion form factor, which is based on Eq. (10), with various options for other contributions, see Sec. III. The effect did not disappear and the value of the ω mass shift was very stable against all changes. Moreover, we have found that the inclusion of the $\phi(1020)$ contribution further improves the quality of the fit not only for our parametrization (where the quality was already excellent), but also very significantly for the GS parametrization, see Tab. VII.

V. SUMMARY AND COMMENTS

The main results of this work are summarized in Tables VIII, IX, and X.

Table VIII shows how well are the particular data sets (CMD-2, SND, KLOE) fitted by two parametrizations (GS and ours) we considered. As a measure of the goodness-of-fit we used the confidence level calculated from the values of χ^2 and the numbers of degrees of freedom shown in Tables III–VII using the CERNLIB routine PROB. The agreement of both the GS and our parametrizations with both the CMD-2 and SND data is on the level that is considered satisfactory in the high-energy physics (based on $\chi^2/NDF \approx 1$). Our parametriza-

tion gives a slightly better confidence level than the GS one, especially for the CMD-2 data.

The disagreement of both parametrizations with the KLOE data is disastrous. But the situation radically improves for our parametrization if we allow the mass of the $\omega(782)$ to vary. Adding this free parameter reduces the original $\chi^2 = 192.6$ to $\chi^2 = 29.6$, what from the statistics point of view fully justifies this step. The confidence level of the hypothesis that our parametrization describes correctly the KLOE data has rocketed to 99.62%. This is highly non-trivial because the statistical errors of the KLOE data are very small, much smaller than those of both sets of data from Novosibirsk. The price that has to be paid for this excellent result is lowering the $\omega(782)$ mass by $\Delta m_\omega = (1.93 \pm 0.19)$ MeV. The physics behind this is unclear. We originally suspected that the measured values of the mass of the two-pion system might contain a systematic shift, at least in the ρ - ω interference region. But the correspondence with the members of the KLOE collaboration excluded this possibility.

Also in the case of the GS parametrization χ^2 decreased from 367.8 to 202.6 when the ω mass was considered an additional free parameter. The required mass shift is even higher than in the case of our parametrization, namely (2.13 ± 0.35) MeV. Adding the ϕ -meson contribution improves the quality of the fit while changing the ω -mass shift negligibly, compare Tabs. VI and VII.

On the other hand, the addition of the ω mass among free parameters is not required by the CMD-2 and SND data. The drop of χ^2 is less than one both for the GS parametrization and ours. Why the KLOE data behaves in a different way is unclear. The issue obviously requires explanation. We do not have any at the moment.

Table IX summarizes the results on the $\rho(770)$ mass we have got by fitting various data sets using the GS and our parametrization. Its salient feature is that for all data sets we considered (CMD-2, SND, KLOE, KLOE with the shifted ω mass) our parametrization gives a smaller $\rho(770)$ mass than the GS one. The differences are large, they range from 7.8 MeV for the KLOE data to 10.3 MeV for the CMD-2 data. At the same time the errors of the masses themselves are very small. It looks like the two columns concerned different quantities. And this is probably the core of the problem. The ρ -meson mass is defined in both parametrizations as a parameter in the corresponding formula for the ρ^0 propagator. As the formulas are different, so must be the definitions and the “experimental” values obtained by fitting those two parametrizations.

The cure of this problem is not easy. To get the “universal” definition of the ρ -meson mass it is insufficient to declare some formula for the propagator (e.g., that based on the GS parametrization) as the canonical one. If that preferred formula is not correct, then the different processes or even the same process measured in different kinematic regions will give different values of the ρ meson mass even if the experiments are ideal. In fact, in order to compare the theoretical predictions with the data on

TABLE IX: The ρ meson mass obtained from various data sets using the GS and our parametrizations

	GS	ours
CMD-2	775.3 ± 1.1	767.08 ± 0.83
SND	774.60 ± 0.52	764.23 ± 0.43
KLOE	769.22 ± 0.32	761.39 ± 0.25
KLOE ^a	771.58 ± 0.60	763.44 ± 0.33
KLOE ^b	772.79 ± 0.35	763.48 ± 0.33

^aAccounting for the energy shift in the ρ - ω region

^bAccounting for the energy shift in the ρ - ω region, $\phi(1020)$ included

TABLE X: The ρ meson width obtained from various data sets using the GS and our parametrizations

Γ_ρ	GS	ours
CMD-2	143.2 ± 2.5	136.1 ± 2.2
SND	147.3 ± 1.0	140.29 ± 0.85
KLOE	144.71 ± 0.56	140.07 ± 0.66
KLOE ^a	146.39 ± 0.51	142.56 ± 0.41
KLOE ^b	151.53 ± 0.55	142.78 ± 0.46

^aFit with the energy shift in the $\rho\omega$ region

^bFit with the energy shift in the $\rho\omega$ region, ϕ contribution included

some process, the propagator has to be convoluted with other factors (“vertices”) that are characteristic for that process and depend also on the virtual ρ -meson momentum squared. If the propagator is not correct, then the cross section or decay width formulas of various processes are incorrect, but each in a different way. We illustrate this point on a toy example in the Appendix.

It is generally believed that the only meaningful definition of the mass and width of a resonance, which does not have asymptotic states, is that in terms of the position of the pole on the second Riemann sheet of the propagator [56]. The pole of the propagator induces a pole in the scattering amplitude (at least in perturbative field theory). The same propagator enters the amplitudes of various processes, so the pole definition should be process independent. The pole position does not change if an analytic function without pole is added to the amplitude to describe the background present in experiments. So the pole definition should be also background independent. In the case of resonances for which a reliable theoretical framework exists ($W^\pm$ and Z^0 bosons) is it possible to relate the mass and width defined in terms of the pole position to the “mass” and “width” that enter as parameters of various fitting formulas [57–59]. The pole-defined mass and width are genuine physical quantities (gauge invariant and renormalization scheme independent in the Z^0 boson case).

There have been several attempts to find the position of the ρ^0 -meson pole on the second sheet of the complex s -plane.

Lang and Mas-Parareda [60] first mapped the cut s -plane into a unit disc with data about the P -wave partial amplitude of the $\pi\pi$ scattering falling on the unit circle. Then they fitted the data to the product of a pole outside the unit circle (corresponding to the second-sheet pole in the s variable) and an analytic function represented by its Taylor series. The pole position and the coefficients in the Taylor series were parameters of the fit. There are different ways to suppress the importance of the higher Taylor series coefficients and make the series finite. Unfortunately, this brings some model dependence. In [60] the Pietarinen [61] smoothness condition was utilized. Lang and Mas-Parareda used five different data sets and got the mutually compatible values of the ρ mass and width, whereas the values of those quantities determined in experimental papers were incompatible. The pole-defined masses were always lower than the masses from the parametrizations used by experimentalists.

Bohacik and Kühnelt [62] used the pole-searching method developed in [63] on the basis of the statistical method for the testing of analyticity [64]. The central objects of the latter are the moments Q_n (n natural) calculated from the data and a smooth function depending on the data errors and density. The moments should obey the $N(0, 1)$ distribution if the investigated data describe an analytic function. If a pole is present, the mean values of the moments are expressed in terms of the pole position and residuum. The authors of [62] mapped the second sheet on the unit disc, so the assumed pole lay inside the unit circle. They either determined the pole parameters from the moments calculated directly from the data or multiplied the data first by the Blaschke pole-killing factor and determined its parameters by requiring the calculated Q_n momenta be $N(0, 1)$ distributed. The resulting mass and width of the ρ meson were compatible with those of Ref. [60]. This approach suffers from model dependence, which crawls in through the construction of the error function and the choice how many and which moments are taken into consideration [64].

Dubnicka and collaborators [23, 29] constructed two analytic models incorporating all pion form factor properties (including pairs of complex conjugate poles on unphysical sheets, corresponding to the ρ , ρ' , and ρ'' resonances) and fitted them to all accessible data. In both models, the (pole) mass of the ρ meson is much smaller than the values given by PDG [19]. The authors tried to keep the number of free parameters as low as possible. As a consequence, the quality of their fits is not very good, $\chi^2/NDF = 1.81$ and 1.58 in [23] and [29], respectively.

The present work has not been intended as a final word in the field of the pion form factor parametrization and extracting the resonance parameters. We are even unable to offer unambiguous values of m_ρ and Γ_ρ . Natural candidates would be those coming from the best fit, which is our fit to the KLOE data. But those are hampered by the presence of a mysterious energy shift in the ρ - ω mass region. The main aim of this paper was to issue the warn-

TABLE XI: Each group fits its data with its own formula

Group	α	R	χ^2
A	0.95	1.096 ± 0.016	11.86
B	1.05	0.866 ± 0.013	10.26
C	0.90	1.369 ± 0.020	4.88

ing that the question of the correct parametrizations is not simple and requires much attention. The resonance parameters obtained by fitting the data are subject of large systematic errors of the parametrizations and have often been underestimated so far.

Acknowledgments

We are indebted to Achim Denig and Graziano Venanzoni from the KLOE collaboration for the enlightening correspondence. One of us (P. L.) acknowledges useful discussions with David Kraus. This work was supported by the Czech Ministry of Education, Youth and Sports under contracts Nos. MSM6840770029 and MSM4781305903 and by the Bergen Computational Physics Laboratory in the framework of the European Community - Access to Research Infrastructure action of the Improving Human Potential Programme. The paper was finished during the visit of one of us (P. L.) to the McGill University. The support from its Physics Department is gratefully acknowledged.

APPENDIX: A TOY EXAMPLE

Let us assume that three experimental groups A, B, and C measured the dependence of the voltage on the electric current of the same wire, but in different intervals. Each interval contained twelve data points. We simulated their measurements assuming the validity of the Ohm law with $R = 1\Omega$ and the Gaussian distribution of experimental values around the Ohm-law values with

the relative error of 5%. The three generated data sets can be accessed at [65]. Each group fitted its data by its own formula. The formulas they chose can be written in a unified way as

$$U = RI^\alpha, \quad (\text{A.1})$$

with $\alpha = 0.95, 1.05$, and 0.90 for the group A, B, and C, respectively. The results of the fits are shown in Table XI. The quality of all fits is acceptable according to the high-energy physics standards and the resulting parameters R have rather small errors. But the R s of the different groups are incompatible.

Then the groups realized that problem lies in different formulas they used and agreed to adopt the formula of the B group ($\alpha = 1.05$) as the common one. They again got the good fits and better agreement among their R

TABLE XII: Each group fits its data with the common formula ($\alpha = 1.05$)

Group	R	χ^2
A	0.928 ± 0.013	10.47
B	0.866 ± 0.013	10.26
C	0.854 ± 0.012	6.10

TABLE XIII: Each group fits its data with the correct formula ($\alpha = 1$)

Group	R	χ^2
A	1.011 ± 0.015	4.64
B	0.987 ± 0.014	8.12
C	0.999 ± 0.014	5.18

parameters, but some inconsistency, especially between groups A and C, remained, see Table XII.

Only after the groups learnt about the correct formula and decided to use common $\alpha = 1$, they have obtained both good fits and compatible results, as it is witnessed in Table XIII.

-
- [1] H. W. Barz *et al.*, Phys. Lett. B **265**, 219 (1991).
 - [2] V. L. Eletsky, B. L. Ioffe, and J. I. Kapusta, Eur. Phys. J. A **3**, 381 (1998).
 - [3] R. Rapp and J. Wambach, Adv. Nucl. Phys. **25**, 1 (2000).
 - [4] G. E. Brown and M. Rho, Phys. Rep. **363**, 85 (2002).
 - [5] O. Teodorescu, A. K. Dutt-Mazumder, and C. Gale, Phys. Rev. C **66**, 015209 (2002).
 - [6] P. F. Kolb and M. Prakash, Phys. Rev. C **67**, 044902 (2003).
 - [7] S. Pratt and W. Bauer, Phys. Rev. C **68**, 064905 (2003).
 - [8] E. V. Shuryak and G. E. Brown, Nucl. Phys. A **717**, 322 (2003).
 - [9] R. Rapp, Nucl. Phys. A **725**, 254 (2003).
 - [10] G. E. Brown and M. Rho, Phys. Rept. **396**, 1 (2004).
 - [11] J. Ruppert, T. Renk, and B. Müller, Phys. Rev. C **73**, 034907 (2006).
 - [12] K. Hikasa *et al.* (Particle Data Group), Phys. Rev. D **45**, S1 (1992).
 - [13] L. Montanet *et al.* (Particle Data Group), Phys. Rev. D **50**, 1173 (1994).
 - [14] R. M. Barnett *et al.* (Particle Data Group), Phys. Rev. D **54**, 1 (1996).
 - [15] C. Caso *et al.* (Particle Data Group), Eur. Phys. J. C **3**, 1 (1998).
 - [16] D. E. Groom *et al.* (Particle Data Group), Eur. Phys. J. C **15**, 1 (2000).

- [17] K. Hagiwara *et al.* (Particle Data Group), Phys. Rev. D **66**, 010001 (2002).
- [18] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
- [19] W.-M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
- [20] M. F. Heyn and C. B. Lang, Z. Phys. C **7**, 169 (1981).
- [21] B. V. Geshkenbein and M. V. Terentev, Yad. Fiz. **40**, 758 (1984) [Sov. J. Nucl. Phys. **40**, 487 (1984)].
- [22] B. V. Geshkenbein, Z. Phys. C **45**, 351 (1989).
- [23] S. Dubnička and L. Martinovič, J. Phys. G **15**, 1349 (1989).
- [24] A. Bernicha, G. López Castro, and J. Pestieau, Phys. Rev. D **50**, 4454 (1994).
- [25] H. B. O'Connell, A. W. Thomas, and A. G. Williams, Nucl. Phys. A **623**, 559 (1997).
- [26] S. Gardner and H. B. O'Connell, Phys. Rev. D **57**, 2716 (1998); Erratum *ibid.* **62**, 019903 (2000).
- [27] M. Benayoun *et al.*, Eur. Phys. J. C **2**, 269 (1998).
- [28] M. Benayoun, H. B. O'Connell, and A. G. Williams, Phys. Rev. D **59**, 074020 (1999).
- [29] S. Dubnička, A. Z. Dubníčková, and A. Antušek, Acta Phys. Slovaca **49**, 165 (1999).
- [30] D. Melikhov, O. Nachtmann, V. Nikonov, and T. Paulus, Eur. Phys. J. C **34**, 345 (2004).
- [31] G. J. Gounaris and J. J. Sakurai, Phys. Rev. Lett. **21**, 244 (1968).
- [32] J. Pišút and M. Roos, Nucl. Phys. B **6**, 325 (1968).
- [33] V. L. Auslender *et al.*, Phys. Lett. B **25**, 433 (1967).
- [34] M. Roos, Nucl. Phys. B **97**, 165 (1975).
- [35] M. Feuillat, J. L. Lucio M, and J. Pestieau, Phys. Lett. B **501**, 37 (2001).
- [36] R. R. Akhmetshin *et al.*, Phys. Lett. B **527**, 161 (2002).
- [37] R. R. Akhmetshin *et al.*, Phys. Lett. B **578**, 285 (2004).
- [38] P. Lichard, Phys. Rev. D **60**, 053007 (1999).
- [39] M. N. Achasov *et al.*, hep-ex/0605013.
- [40] A. Aloisio *et al.*, Phys. Lett. B **606**, 12 (2005).
- [41] G. R. Farrar and D. R. Jackson, Phys. Rev. Lett. **43**, 246 (1979); G. P. Lepage and S. J. Brodsky, Phys. Lett. B **87**, 359 (1979); A. V. Efremov and A. V. Radyushkin, Phys. Lett. B **94**, 245 (1980).
- [42] G. F. Chew and S. Mandelstam, Phys. Rev. **119**, 467 (1960).
- [43] P. Federbush, M. L. Goldberger, and S. B. Treiman, Phys. Rev. **112**, 642 (1958); S. Gasiorowicz, *Elementary Particle Physics* (Wiley, New York, 1966).
- [44] M. T. Vaughn and K. C. Wali, Phys. Rev. Lett. **21**, 938 (1968).
- [45] R. Barate *et al.* (ALEPH Collaboration), Z. Phys. C **76**, 15 (1997).
- [46] V. M. Aulchenko *et al.*, Pisma v Zh. Eksp. Teor. Fiz. **82**, 841 (2005) [JETP Lett. **82**, 743 (2005)].
- [47] R. Kokoski and N. Isgur, Phys. Rev. D **35**, 907 (1987).
- [48] N. Isgur, C. Morningstar, and C. Reader, Phys. Rev. D **39**, 1357 (1989).
- [49] M. Conversi *et al.*, Phys. Lett. B **52**, 493 (1974); B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **71**, 052001 (2005).
- [50] F. James and M. Roos, Comput. Phys. Commun. **10**, 343 (1975).
- [51] L. M. Barkov *et al.*, Nucl. Phys. B **256**, 365 (1985).
- [52] S. R. Amendolia *et al.*, Nucl. Phys. B **277**, 168 (1986).
- [53] M. N. Achasov *et al.*, Zh. Eksp. Teor. Fiz. **128**, 1201 (2005) [J. Exp. Theor. Phys. **101**, 1053 (2005)].
- [54] Min-Shih Chen and P. M. Zerwas, Phys. Rev. D **11**, 58 (1975); H. Czyż, hep-ph/0606227.
- [55] A. Denig and G. Venanzoni, private communication.
- [56] R. E. Peierls, Proceedings of Glasgow Conference on Nuclear and Meson Physics (Pergamon Press, Inc., London, 1954), p. 296; P. T. Matthews and A. Salam, Phys. Rev. **112**, 283 (1958); **115**, 1079 (1959); M. Lévy, Nuovo Cimento **13**, 115 (1959); J. Schwinger, Ann. Phys. **9**, 169 (1960); R. Jacob and R. G. Sachs, Phys. Rev. **121**, 350 (1961).
- [57] S. Willenbrock and G. Valencia, Phys. Lett. B **259**, 373 (1991).
- [58] R. G. Stuart, Phys. Lett. B **262**, 113 (1991).
- [59] A. Sirlin, Phys. Rev. Lett. **67**, 2127 (1991).
- [60] C. B. Lang and A. Mas-Parareda, Phys. Rev. D **19**, 956 (1979).
- [61] E. Pietarinen, Nuovo Cimento A **12**, 522 (1972); Nucl. Phys. B **55**, 541 (1973).
- [62] J. Bohacik and H. Kühnelt, Phys. Rev. D **21**, 1342 (1980).
- [63] A. Nogová, J. Pišút, and P. Prešnajder, Nucl. Phys. B **61**, 438 (1973).
- [64] A. Pázman *et al.*, Nucl. Phys. B **46**, 637 (1972); P. Lichard, J. Pišút, and P. Prešnajder, *Representation of Data by Analytic Functions in Physics* (Veda, Bratislava, 1983) (in Slovak).
- [65] <http://axpsu.fpf.slu.cz/~lic10uf/0hm.html>.